

Coordinate Changes for Integrals

In Calc I we solved things like

$$\int_0^5 x e^{x^2} dx \quad \left(\begin{array}{l} u = x^2 \\ du = 2x dx \end{array} \right) \leftarrow \text{parameter change}$$

In double integrals, we made a polar coordinate change

$$dA_{\text{cart}} = r dA_{\text{polar}}$$

Q: How do we deal with more general changes?

A: We use the Jacobian to understand diff. changes

Def: Suppose $\begin{cases} x_1 = x_1(u_1, u_2, \dots, u_n) \\ x_2 = x_2(u_1, u_2, \dots, u_n) \\ \vdots \\ x_n = x_n(u_1, u_2, \dots, u_n) \end{cases}$ is a

coordinate change by diff functions. The Jacobian of the coordinate change is

$$\frac{\partial(x_1, x_2, \dots, x_n)}{\partial(u_1, u_2, \dots, u_n)} = \begin{vmatrix} \frac{\partial x_1}{\partial u_1} & \frac{\partial x_1}{\partial u_2} & \dots & \frac{\partial x_1}{\partial u_n} \\ \frac{\partial x_2}{\partial u_1} & \frac{\partial x_2}{\partial u_2} & \dots & \frac{\partial x_2}{\partial u_n} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial x_n}{\partial u_1} & \frac{\partial x_n}{\partial u_2} & \dots & \frac{\partial x_n}{\partial u_n} \end{vmatrix} \det$$

Ex: Compute the signed Jacobian of Polar transform $\begin{cases} x = r \cos(\theta) \\ y = r \sin(\theta) \end{cases}$

$$\begin{aligned} \text{Sol: } \frac{\partial(x, y)}{\partial(r, \theta)} &= \det \begin{bmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} \end{bmatrix} = \det \begin{bmatrix} \cos \theta & -r \sin \theta \\ \sin \theta & r \cos \theta \end{bmatrix} \\ &= \cos \theta \cdot r \cos \theta - (-r \sin \theta \sin \theta) = r(\cos^2 \theta + \sin^2 \theta) = r \end{aligned}$$

NB: swapping the order

$$\begin{aligned} \frac{\partial(x, y)}{\partial(\theta, r)} &= \det \begin{bmatrix} -r \sin \theta & \cos \theta \\ \sin \theta & r \cos \theta \end{bmatrix} \\ &= -r \sin^2 \theta - r \cos^2 \theta = -r \end{aligned}$$

The unsigned Jacobian is $\left| \frac{\partial(x_1, x_2, \dots, x_n)}{\partial(u_1, u_2, \dots, u_n)} \right|$

Prop: If $f(x_1, x_2, \dots, x_n)$ is a cts function and

$$\begin{cases} x_1 = x_1(u_1, u_2, \dots, u_n) \\ x_2 = x_2(u_1, u_2, \dots, u_n) \\ \vdots \\ x_n = x_n(u_1, u_2, \dots, u_n) \end{cases} \text{ is a diff coord trans}$$

$$\int_{R_{\text{old}}} f dV_{\text{old}} = \int_{R_{\text{old}}} f(x_1(u_1, \dots, u_n), \dots, (x_n(u_1, \dots, u_n)) \left| \frac{\partial(x_1, \dots, x_n)}{\partial(u_1, \dots, u_n)} \right|$$

NB: This matches with our work for polar coords

Example: Compute $\iint_R (x-3y) dA$ for R the triangle

with vertices $(0,0), (1,2), (2,1)$

Sol: is the parameterization when



$$(\alpha, \beta) = (1,0) \text{ and } (x(\alpha, \beta), y(\alpha, \beta)) = (1,2)$$

$$(2,1) \text{ and when } (\alpha, \beta) = (0,1)$$

$$\text{we have } (x(\alpha, \beta), y(\alpha, \beta)) = (1,2)$$

$$\text{and when } (\alpha, \beta) = (0,0) \text{ yields}$$

$$(x(\alpha, \beta), y(\alpha, \beta)) = (0,0)$$



By high school geometry, this linear

change takes R_{new} to R_{old}

$$R_{\text{new}} = \{(\alpha, \beta) : 0 \leq \alpha \leq 1, 0 \leq \beta \leq 1-\alpha\}$$

$$\text{The Jacobian is } \frac{\partial(x,y)}{\partial(\alpha,\beta)} = \det \begin{bmatrix} \frac{\partial x}{\partial \alpha} & \frac{\partial x}{\partial \beta} \\ \frac{\partial y}{\partial \alpha} & \frac{\partial y}{\partial \beta} \end{bmatrix} = \det \begin{bmatrix} 2 & 1 \\ 1 & 2 \end{bmatrix} = 3$$

$$\therefore \iint_{R_{\text{old}}} (x-3y) dA_{\text{old}} = \iint_{R_{\text{new}}} ((2\alpha+\beta) - 3(\alpha+2\beta)) \left| \frac{\partial(x,y)}{\partial(\alpha,\beta)} \right| dA_{\text{new}}$$

$$= \int_{\alpha=0}^1 \int_{\beta=0}^{1-\alpha} (-\alpha-5\beta) 3 d\beta d\alpha$$

$$= -3 \int_0^1 \int_0^{1-\alpha} (\alpha+5\beta) d\beta d\alpha$$

$$= -3 \int_0^1 \left[\alpha\beta + \frac{5}{2}\beta^2 \right]_0^{1-\alpha} d\alpha$$

$$= -3 \int_0^1 (1-\alpha) \left(\alpha + \frac{5}{2}(1-\alpha) \right) d\alpha$$

$$= -\frac{3}{2} \int_0^1 (5\alpha - 4\alpha^2 + \alpha^3) d\alpha = -3$$

Now let's generalize polar coords in 3-space

I. Naive way. Cylindrical coords

IDEA: parameterize the plane by polar coords

leave the orthogonal axis unchanged

$$\text{Ex: } \begin{cases} x = r \cos \theta \\ y = r \sin \theta \\ z = z \end{cases}$$



Jacobian is

$$J(x, y, z) = \det \begin{bmatrix} \frac{\partial x}{\partial r} & \frac{\partial x}{\partial \theta} & \frac{\partial x}{\partial z} \\ \frac{\partial y}{\partial r} & \frac{\partial y}{\partial \theta} & \frac{\partial y}{\partial z} \\ \frac{\partial z}{\partial r} & \frac{\partial z}{\partial \theta} & \frac{\partial z}{\partial z} \end{bmatrix} = \det \begin{bmatrix} \cos \theta & -r \sin \theta & 0 \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$= \cos \theta (r \cos \theta \cdot 1 - 0) - (-r \sin \theta) (\sin \theta \cdot 1 - 0) + 0 (\sin \theta \cdot 0 - r \cos \theta \cdot 0) \\ = r \cos^2 \theta + r \sin^2 \theta = r$$

∴ when we compute an integral in cylindrical coords,
we need to multiply the differential by r
* This is true of all cylindrical changes

Ex: compute $\iiint_R (x+y+z) dV$ for the solid with first octant and below paraboloid $4-x^2-y^2=z$

Sol: In coords $\begin{cases} x = r \cos \theta \\ y = r \sin \theta \\ z = z \end{cases}$



$$\Rightarrow R_{\text{cyl}} = \{(r, \theta, z) : 0 \leq \theta \leq \frac{\pi}{2}, 0 \leq r \leq 2, 0 \leq z \leq 4-r^2\}$$

$$\begin{aligned} \therefore \iiint_R (x+y+z) dV &= \int_{\theta=0}^{\frac{\pi}{2}} \int_{r=0}^2 \int_{z=0}^{4-r^2} (r \cos \theta + r \sin \theta + z) r dz dr d\theta \\ &= \int_0^{\frac{\pi}{2}} \int_0^2 r [r \sin \theta - r \cos \theta + z\theta]_0^{4-r^2} dz dr \\ &= \int_0^{\frac{\pi}{2}} \int_0^2 r (1-r^2 + z\frac{\pi}{2}) - (0-r+0) dz dr \\ &= \int_0^{\frac{\pi}{2}} \int_0^2 (2r^2 + \frac{\pi}{2} z r) dz dr \\ &= \int_0^{\frac{\pi}{2}} [2r^2 z + \frac{\pi}{4} z^2 r]_{z=0}^{4-r^2} dr \\ &= \int_0^{\frac{\pi}{2}} (2r^2(4-r^2) + \frac{\pi}{4} r(4-r^2)^2) dr \\ &= \int_0^{\frac{\pi}{2}} (8r^2 - 2r^4 + \frac{\pi}{4} (16r - 8r^3 + r^5)) dr \\ &= \left[\frac{8}{3} r^3 - \frac{2}{5} r^5 + \frac{\pi}{4} (8r^2 - 2r^4 + \frac{1}{6} r^6) \right]_0^{\frac{\pi}{2}} \\ &= \frac{64}{3} - \frac{32}{5} + \frac{\pi}{4} (32 - 32 + \frac{32}{3}) - 0 \\ &= \frac{64}{3} - \frac{32}{5} + \frac{8\pi}{3} \end{aligned}$$

II: Spherical Coordinates

In spherical coordinates, we parameterize points (x, y, z) using 3 pieces of data

ρ = distance from origin

θ = angle made w/ +x-axis and point $(x, y, 0)$

ϕ = angle made w/ +z-axis and point (x, y, z)

Note: $\sin(\phi) = \frac{r}{\rho}$, so $r = \rho \sin(\phi)$

\therefore in our parameterization

$$\begin{cases} x = r \cos(\theta) = \rho \sin(\phi) \cos(\theta) \\ y = r \sin(\theta) = \rho \sin(\phi) \sin(\theta) \end{cases}$$

Also, $\cos(\phi) = \frac{z}{\rho}$, so $z = \rho \cos(\phi)$

Hence the spherical coord parameterization is

$$\begin{cases} x = \rho \sin(\phi) \cos(\theta) \\ y = \rho \sin(\phi) \sin(\theta) \\ z = \rho \cos(\phi) \end{cases}$$

